

Fermions as $U(1)$ instantons

M.N. Stoilov

*Bulgarian Academy of Sciences,
Institute of Nuclear Research and Nuclear Energy,
Blvd. Tzarigradsko Chaussee 72, Sofia 1784, Bulgaria*
e-mail: mstoilov@inrne.bas.bg

February 2, 2008

Abstract

Anomalous quantization of the electromagnetic field allows non-trivial (anti) self-dual configurations to exist in four-dimensional Euclidian space-time. These instanton-like objects are described as massless spinor particles.

PACS: 12.20.-m, 14.80.Hv

key words: Electrodynamics, instanton, anomalous quantization

Quantization in the vicinity of a fixed field configuration which is a solution of the equations of motion is a common technique in Quantum Field Theory. The classical configuration around which we quantize represents the vacuum of the model and the deviation of the field from this vacuum is the quantum field. When the classical solution/vacuum is not unique but depends on some parameters then a gauge symmetry emerges in the model [1]–[3]. In this case the standard path integral over all fields is replaced by functional integral (with suitable measure which takes into account the gauge symmetry) over the quantum field plus ordinary integrals over all parameters on which the vacuum configuration depends [1]. Here we shall consider an example where the vacuum is again not unique but now we do not know its explicit functional form. Instead of this we know that it satisfies some well defined conditions which do not coincide with the equations of motion. Our example is pure Electrodynamics in four-dimensional Euclidian space-time E^4 and the classical solution around which we decompose the electromagnetic potential is (anti) self-dual. This seems rather trivial because it is known that for simple topological reasons there are no instanton configurations in Electrodynamics. However, the conditions for (anti) self-duality can be rewritten as conditions for Fueter quaternion (anti) analyticity. It is shown in Ref.[4] that fields which are Fueter (anti) analytic can be quantized either as fermions or as bosons. We use this result to quantize anomalously the (anti) self-dual electromagnetic field. This is the key moment in the work — we use spinors to describe instantons. The change of the statistics ‘stabilizes’ the instantons allowing non-trivial (anti) self-dual configurations to exist. The idea that instantons change the statistics of some fields is not new in the non-Abelian case [5]–[7]. Here these fields are instantons themselves. The topological reason for the existence of $U(1)$ instantons is that using spinors we effectively change the base of the $U(1)$ bundle from E^4 to its double covering space which has non-trivial fundamental group. Our conclusion is that anomalously quantized instanton modes of the electromagnetic field are described by the fermion field in massless spinor Electrodynamics.

The use of a non-unique classical solution as vacuum requires some precautions. In this case the quantum evolution operator which determines the propagator of the quantum field (see below for the exact definition) has zero modes. This means that there is gauge symmetry in the resulting model, with zero modes describing Goldstone bosons. The existence of a zero mode is rather easy to demonstrate. Suppose we have a model for a field φ with Lagrangean L and let φ_{cl} is a classical solution of the equations of motion, i.e.,

$$\frac{\partial L}{\partial \varphi}|_{\varphi=\varphi_{cl}} = 0. \quad (1)$$

Let us expand the field φ around the classical solution φ_{cl}

$$\varphi = \varphi_{cl} + \eta. \quad (2)$$

We consider the field η as a small quantum variation of the field φ (quantum field) and φ_{cl} as vacuum. Putting the expansion (2) for φ into the Lagrangean L and keeping terms up to second order with respect to η we get

$$L'(\varphi_{cl}, \eta) = L(\varphi_{cl}) + \frac{1}{2} \left(\frac{\partial^2 L}{\partial \varphi \partial \varphi} \Big|_{\varphi=\varphi_{cl}} \right) \eta^2. \quad (3)$$

This truncated Lagrangean is considered as a Lagrangean for the independent field η . The operator $T = \frac{1}{2}(\partial^2 L / \partial \varphi \partial \varphi)|_{\varphi=\varphi_{cl}}$ determines the propagator of the quantum field. This is the operator which is called ‘quantum evolution operator’. Now suppose that φ_{cl} depends on some parameter, say α , and let us calculate the α -derivative of the equation of motion (1). The result is

$$0 = \frac{\partial}{\partial \alpha} \left(\frac{\partial L}{\partial \varphi} \Big|_{\varphi=\varphi_{cl}} \right) = T \frac{\partial \varphi_{cl}}{\partial \alpha}. \quad (4)$$

Therefore, $\partial \varphi_{cl} / \partial \alpha$ is a zero mode of the quantum evolution operator or in other words, there is a gauge symmetry in the Lagrangean L' . This gauge symmetry emerges no matter what the initial Lagrangean L is. The symmetry has to be taken into account when we write down the transition amplitude for the model with Lagrangean (3). For this model the functional integration over the field φ transforms into integral over α and functional integral over η

$$\int D\varphi e^{-\int dt L} \rightarrow \int d\alpha \int D\eta e^{-\int dt L'} \delta(\chi) \Delta \quad (5)$$

where χ is some gauge fixing term and Δ is the corresponding Faddeev–Popov determinant. The integration over the parameter α effectively sums different classical solutions (vacuums) in the model.

It has to be stressed that it does not follow automatically from the above considerations that when the Lagrangean L possesses *ad initium* gauge symmetry then L' has bigger gauge symmetry. The Lagrangean (3) can have the same symmetry as L if $\partial \varphi_{cl} / \partial \alpha$ coincides with some of the zero modes due to the initial gauge freedom. The gauge symmetry of L' could be even smaller than the symmetry of L if φ_{cl} is gauge non-invariant. In any case one has to determine exactly the symmetry of the quantum evolution operator and to use adequate gauge conditions.

In the present work we consider a model for which we do not know the explicit form of φ_{cl} and the parameters which it depends on, but we know

that the field configuration we use as vacuum satisfies the equation $M\varphi_{cl} = 0$, where M is some operator. In this case eq.(5) takes the form

$$\int D\varphi e^{-\int dtL} \rightarrow \int D\varphi_{cl} \delta(M\varphi_{cl}) D\eta e^{-\int dtL'} \delta(\chi) \Delta. \quad (6)$$

The integration over α in (5) is replaced in (6) by $\int D\varphi_{cl} \delta(M\varphi_{cl})$ which is the way to sum over all possible vacuums in the present case. As a result now φ_{cl} and η describe two different fields. The propagator of η is determined as above by T and the propagator of φ_{cl} is determined by $L(\varphi_{cl})$ and $\log(\det(M))$.

The model we shall deal with hereafter is pure Electrodynamics in four-dimensional Euclidian space-time E^4 . The model is defined by the following action

$$\mathbf{A} = - \int \frac{1}{4} F_{\mu\nu} F_{\mu\nu}. \quad (7)$$

Here $F_{\mu\nu}$ is the field strength tensor for the electromagnetic potential A_μ ; $F_{\mu\nu} = \partial_\nu A_\mu - \partial_\mu A_\nu$. In four-dimensional space-time it is possible to use quaternions to describe the electromagnetic field [8], [9]. The use of quaternions is especially convenient in E^4 .

Let us remind the definition of quaternion number $\mathcal{Q}$

$$\mathcal{Q} = q_0 + q_i e_i \quad (8)$$

where q_μ , $\mu = 0, \dots, 3$ are four real numbers and e_i , $i = 1, 2, 3$ are three non-commutative imaginary units such that:

$$e_i e_j = -\delta_{ij} + \epsilon_{ijk} e_k. \quad (9)$$

Here ϵ_{ijk} is the totally anti symmetric third rank tensor. There is a natural operation of conjugation for quaternions. The quaternion which is conjugated to $\mathcal{Q}$ we shall denote by $\bar{\mathcal{Q}}$ where

$$\bar{\mathcal{Q}} = q_0 - q_i e_i. \quad (10)$$

For quaternion functions there is a notion analogous to analyticity for complex functions — the so called Fueter analyticity. Let us define the first order differential operators $\bar{\mathcal{D}}$ and $\mathcal{D}$

$$\bar{\mathcal{D}} = \partial_0 + e_i \partial_i \quad (11)$$

$$\mathcal{D} = \partial_0 - e_i \partial_i. \quad (12)$$

The function $\mathcal{F}$ is called Fueter analytic if it satisfies the following equation

$$\bar{\mathcal{D}}\mathcal{F} = 0. \quad (13)$$

If the function satisfies the equation

$$\mathcal{D}\mathcal{F} = 0 \quad (14)$$

then the function is Fueter anti analytic.

Having the electromagnetic potential A_μ we can associate two quaternion functions $\mathcal{A}$ and $\bar{\mathcal{A}}$ to it

$$\mathcal{A} = A_0 + A_i e_i \quad (15)$$

$$\bar{\mathcal{A}} = A_0 - A_i e_i. \quad (16)$$

There is also a possibility [8] to associate a complex-valued, imaginary quaternion function directly to the electromagnetic field strength tensor $F_{\mu\nu}$. For this function it is shown in the same paper that the free Maxwell equations are conditions for Fueter analyticity. Having in mind this result we want to see what the Fueter analyticity condition gives for the electromagnetic potential. Using eqs.(9, 11, 12, 15, 16) we get the following result

$$\mathcal{D}\mathcal{A} = \partial_\mu A_\mu + (\partial_0 A_i - \partial_i A_0 - \epsilon_{ijk} \partial_j A_k) e_i \quad (17)$$

$$\bar{\mathcal{D}}\bar{\mathcal{A}} = \partial_\mu A_\mu + (-\partial_0 A_i + \partial_i A_0 - \epsilon_{ijk} \partial_j A_k) e_i \quad (18)$$

So, if the electromagnetic potential is such that

$$\mathcal{D}\mathcal{A} = 0 \quad (19)$$

then we have self-dual configuration plus Lorentz gauge condition. On the other hand if

$$\bar{\mathcal{D}}\bar{\mathcal{A}} = 0 \quad (20)$$

then the field is anti self-dual plus again the Lorentz condition.

In perturbation theory the propagator of the electromagnetic field is Green's function of the second order Maxwell's equations. This Green's function in momentum space behaves as $1/p^2$. On the other hand, the instantons satisfy first order differential equations and so their propagator in momentum space looks like $1/p$. Therefore perturbation theory does not take into account the propagation of (anti) self-dual configurations. In order to take into account the instanton propagation we expand formally the electromagnetic potential A around a vacuum which is a sum of fixed self-dual potential A^+ and anti self-dual potential A^- , i.e. we write

$$A_\mu = A_\mu^+ + A_\mu^- + A'_\mu. \quad (21)$$

Then we allow $A^\pm$ to vary still remaining (anti) self-dual and using formula (6) we find the path integral measure for the instantons. This allows us to

determine their propagator. Carrying out this strategy we first introduce the field strengths F^+ , F^- and F' which correspond to the potentials A^+ , A^- and A' respectively. The sum $A^+ + A^-$ is a classical solution because both A^+ and A^- are classical solutions and because Maxwell's equations are linear. Using this fact and the following well-known relations

$$\begin{aligned} F_{\mu\nu}^+ F_{\mu\nu}^- &= 0 \\ F_{\mu\nu}^+ F_{\mu\nu}^+ &= 2\partial_\lambda \left(\epsilon_{\lambda\mu\nu\rho} A_\mu^+ \partial_\nu A_\rho^+ \right) \\ F_{\mu\nu}^- F_{\mu\nu}^- &= -2\partial_\lambda \left(\epsilon_{\lambda\mu\nu\rho} A_\mu^- \partial_\nu A_\rho^- \right) \end{aligned}$$

it is easy to see that the electromagnetic action (7) in the vicinity of the field configuration $A^+ + A^-$ takes the form

$$\mathbf{A} = - \int \frac{1}{4} F_{\mu\nu}' F_{\mu\nu}'. \quad (22)$$

As a consequence of eq.(22) the quantum evolution operator in our case coincides with the Maxwell's operator. There are no extra zero modes and the gauge symmetry group is $U(1)$. Certainly, in order to obtain a well-defined transition amplitude, we have to choose a gauge fixing function. Once again, because of eq.(22) the gauge fixing function can be each of the standard ones used in Electrodynamics (now written for the field A'). Therefore we can view A' as the standard electromagnetic potential and eq.(22) as the standard pure electromagnetic action.

Another consequence of eq.(22) is that because $\mathbf{A}$ does not depend on A^+ and A^- the contribution of (anti) instanton modes to the transition amplitude will be a multiplicative constant which gives the number of different (anti) self-dual configurations. In order to find this number we have to clarify what 'different (anti) self-dual configurations' means, which is the question for the gauge freedom in A^+ and A^- . Writing eq.(21) we have assumed that A' is a connection as well as A is. Now, note that the difference of two connections is a tensor (invariant in our $U(1)$ case). So, the proper handling of the expansion (21) requires to use such self-dual and anti self-dual potentials so that their sum $A^+ + A^-$ is gauge invariant, i.e.

$$\delta_\epsilon(A_\mu^+ + A_\mu^-) = 0. \quad (23)$$

Here δ_ϵ denotes the usual $U(1)$ gauge variation whose parameter ϵ is an arbitrary function on E^4 . However, even if $A^+ + A^-$ satisfies eq.(23) there is still room for the following gauge transformation:

$$\delta'_\zeta(A_\mu^\pm) = \pm \partial_\mu \zeta. \quad (24)$$

(ζ is an arbitrary function too.) Thus the decomposition (21) introduces a new gauge freedom in the model which is independent from the initial one. But we do not want it, so we have to fix it ensuring that

$$\delta'_\zeta(A_\mu^+ - A_\mu^-) = 0. \quad (25)$$

So, we have to impose two conditions (23) and (25) on two independent combinations of the potentials A^+ and A^- which is equivalent to fixing the gauge separately in both the self-dual and anti self-dual potentials. We choose as gauge conditions

$$\partial_\mu A_\mu^\pm = 0.$$

Therefore the requirements for Fueter anti-analyticity for $\mathcal{A}^+$ and Fueter analyticity for $\mathcal{A}^-$ perfectly specify A^+ and A^- . As a consequence the integration measure over self-dual configurations which counts these configurations properly must be something like

$$\delta(\mathcal{D}\mathcal{A}^+) \mathcal{D}\mathcal{A}^+ \quad (26)$$

with an analogous expression for the anti self-dual potential.

An important note should be added at this point. The representation of the quaternion imaginary units we are using here is

$$e_k = -i\sigma_k, \quad k = 1, 2, 3 \quad (27)$$

where σ_k are the Pauli matrices. As a result, first, in this representation the formulae (15, 16) can be viewed as a change of variables such that now the electromagnetic potential has not one vector index but two spinor ones (vector transforms in bispinor). Second, because of eqs.(27), any quaternion number $\mathcal{Q}$ is now represented by a 2×2 matrix

$$\mathcal{Q} = q_0 + q_i e_i = \begin{pmatrix} a & -\bar{b} \\ b & \bar{a} \end{pmatrix} \quad (28)$$

where $a = q_0 - iq_3$, $b = q_2 - iq_1$ with $\bar{a}$, $\bar{b}$ denoting the complex conjugates of a and b . Therefore, if for some non-zero two-dimensional normalized vector (in fact, 2D normalized Weyl spinor) v , say $v = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$, we know $\mathcal{Q} \cdot v$, then we know $\mathcal{Q}$ itself because four real numbers q_μ are just enough to determine two complex numbers a and b and vice-versa. The same comments are by no means applicable to $\bar{\mathcal{Q}}$. These arguments have to be taken into account when we write down the integration measure. Using the representation (27) we get in eq.(26) a delta function of a matrix which has to be understood as a product of delta functions of each matrix entry. This gives four complex

conditions for A^+ which exceeds the correct number. In order to get two complex conditions for A^+ we have to pick up a constant spinor v which to multiply $\mathcal{D}\mathcal{A}^+$ on the right and to use $\mathcal{D}\mathcal{A}^+ \cdot v$ as an argument of the delta function in eq.(26). Unfortunately, following this recipe we face a problem which is due to the gauge non-invariance of v : the quantity $\mathcal{D}\mathcal{A}^+ \cdot v$ is neither gauge invariant, nor gauge covariant. So, the price we have to pay in order to have the correct number of constraints is gauge non-invariance of the obtained expression. We shall come back to this question later. Let us define the spinors ψ^+ and ψ^- as follows

$$\begin{aligned}\psi^+ &= \mathcal{A}^+ \cdot v \\ \psi^- &= \bar{\mathcal{A}}^- \cdot u.\end{aligned}\tag{29}$$

Then the transition amplitude for pure Electrodynamics plus instanton contributions and without gauge freedom of the quantum field to be fixed takes the form

$$S = \int D\psi^+ D\psi^- DA' \exp \left\{ -\frac{1}{4} \int F'_{\mu\nu} F'_{\mu\nu} \right\} \delta(\mathcal{D}\psi^+) \delta(\bar{\mathcal{D}}\psi^-).\tag{30}$$

Note that using spinors ψ^+ and ψ^- to describe instantons and anti-instantons we effectively change the trivial $U(1)$ bundle over E^4 to a one whose base B is a double cover of E^4 . The fundamental group of B is non-zero: $\pi(B) = \mathbf{Z}$ which is the reason for nontrivial (anti) self dual configurations to exist. These configurations are, however, spinors and we shall quantize them as Fermi particles, i.e., we suppose that v and u are Grassmann odd quantities. This approach could be viewed as an inverse to the procedure used in Ref.[4] for anomalous quantization of the massless spinor field.

The two delta functions in eq.(30) can be represented as one. We introduce four-dimensional Dirac spinor ψ as direct sum of Weil spinors ψ^+ and ψ^-

$$\psi = \begin{pmatrix} \psi^- \\ \psi^+ \end{pmatrix}\tag{31}$$

and make use of the diagonal γ^5 representation of the 4D Euclidian gamma matrices

$$\gamma_0 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \gamma_i = \begin{pmatrix} 0 & i\sigma_i \\ -i\sigma_i & 0 \end{pmatrix}, \quad \gamma_5 = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}.\tag{32}$$

In this representation the 4D Euclidian Dirac operator has the form

$$\partial_\mu \gamma_\mu = \begin{pmatrix} 0 & \partial_0 + i\partial_k \sigma_k \\ \partial_0 - i\partial_k \sigma_k & 0 \end{pmatrix} = \begin{pmatrix} 0 & \mathcal{D} \\ \bar{\mathcal{D}} & 0 \end{pmatrix}.\tag{33}$$

Note that the operators $\bar{\mathcal{D}}$ and $\mathcal{D}$ which appear in eq.(33) are those defined by eqs.(11, 12) taken in the representation (27). Using eqs.(31, 33) we get:

$$\delta(\mathcal{D}\psi^+)\delta(\bar{\mathcal{D}}\psi^-) = \delta(\partial_\mu\gamma_\mu\psi). \quad (34)$$

With the help of Lagrange multipliers the delta function (34) can be represented as part of the action. Because of the spinor character of the delta function argument and because of hermiticity of the action, the Lagrange multipliers have to form a spinor, conjugated to the spinor ψ [4]. Thus we get the following expression for the transition amplitude:

$$S = \int D\bar{\psi}D\psi DA' \exp \left\{ - \int i\bar{\psi}\partial_\mu\gamma_\mu\psi + \frac{1}{4}F'_{\mu\nu}F'_{\mu\nu} \right\}. \quad (35)$$

It has already been mentioned above that the introduction of the spinors v and u spoils gauge covariance of the delta function argument. As a result the action in eq.(35) is gauge non-invariant. However, we have started with gauge invariant action, and we have to restore this invariance in eq.(35). We do this in the standard way, namely prolonging the derivatives with A' thus obtaining

$$S = \int D\bar{\psi}D\psi DA' \exp \left\{ - \int \bar{\psi}(i\partial_\mu + A'_\mu)\gamma_\mu\psi + \frac{1}{4}F'_{\mu\nu}F'_{\mu\nu} \right\}. \quad (36)$$

Eq.(36) coincides with the expression for the transition amplitude of massless spinor Electrodynamics. In our case the spinor field represents the anomalously quantized instanton modes. Once again, we stress that gauge is not fixed in eq.(30) and so it is not fixed in (36) as well. There is nothing special in this gauge fixing — it is a standard gauge fixing for the electrodynamics potential A' and we shall not specify it explicitly.

It is possible to get eq.(36) directly without passing through eq.(35). For this purpose we have to use in eq.(29) a gauge invariant spinor w instead of the constant but gauge non-invariant spinor v , where w is given by the following expression:

$$w = e^{-i\phi[\Gamma]}v. \quad (37)$$

Here $\phi[\Gamma]$ is a phase which compensates the gauge transformation of v . The phase $\phi[\Gamma]$ is non-integrable [10], i.e. it is not a function with definite value in each space-time point x , but instead it is a multi-valued functional depending on the path Γ along which we reach the point x . However, the phase $\phi[\Gamma]$ possesses well defined derivatives

$$\partial_\mu\phi[\Gamma] = A'_\mu. \quad (38)$$

Formally, $\phi[\Gamma]$ can be written as an integral over path Γ

$$\phi = \int_{\Gamma}^x dx_{\mu} A'_{\mu}(x). \quad (39)$$

When we use w to reduce the number of constraints on A^+ we get

$$\begin{aligned} \mathcal{D}\mathcal{A}^+ \cdot w &= \mathcal{D}(e^{-i\phi[\Gamma]}\mathcal{A}^+ \cdot v) \\ &= e^{-i\phi[\Gamma]}(\mathcal{D} - i\mathcal{A}')\psi^+ \end{aligned} \quad (40)$$

and this gauge invariant expression we have to put into the delta function which defines the measure over self-dual field configurations. Representing the delta function as an exponent, the phase $e^{-i\phi[\Gamma]}$ in eq.(40) will be compensated by the (conjugated to it) phase of the Lagrange multipliers. Thus the phase $\phi[\Gamma]$ does not appear at all in the Lagrangean. Applying the same procedure to the measure which counts anti self-dual fields, we get for the transition amplitude directly the gauge invariant expression (36).

Let us summarize our results: we have started with pure Electrodynamics with vacuum which is (anti) self-dual, then we quantize anomalously the vacuum modes and what we get finally is massless spinor Electrodynamics where fermions describe instanton modes.

Acknowledgement

It is a pleasure to thank E. Nissimov and S. Pacheva for useful discussions and to O. Stoytchev for reading the manuscript.

Dedication

The work is dedicated to the memory of Prof. Dimitar Stoyanov

References

- [1] J.-L. Gervais and B. Sakita, Phys. Rev. **D 10** (1975) 2943.
- [2] J. Goldstone and R. Jackiw, Phys. Rev. **D 11** (1975) 1486.
- [3] E. Tomboulis, Phys. Rev. **D 12** (1975) 1678
- [4] D.Ts. Stoyanov, J. Phys. A: Math. Gen. **25** (1992) 4245.
- [5] R. Jackiw and C. Rebbi, Phys. Rev. **D 13** (1976) 3398.

- [6] P. Hasenfratz and G. 't Hooft, Phys. Rev. Lett. 36 (1976) 1119.
- [7] A. Goldhaber, Phys. Rev. Lett. 36 (1976) 1122.
- [8] D.Ts. Stoyanov, *Quaternion analyticity and Maxwell equations*, in Theoretic and High Energy Physics, BAS, Sofia, 1988.
- [9] see, e.g. Y. Q. Gu, *Canonical Form of Field Equations*, arXiv:hep-th/0610189 and references therein.
- [10] P.A.M. Dirac, Proc. Roy. Soc. **A** 133 (1931) 60.